

GROUP 4 HERBICIDE

ORACLE®
Dicamba Agricultural Herbicide

COMMERCIAL

GUARANTEE:

Dicamba..... 480 g/L
(Present as Dimethylamine Salt)

REGISTRATION NUMBER 26722 PEST CONTROL PRODUCTS ACT

READ THE LABEL AND BOOKLET BEFORE USING

KEEP OUT OF REACH OF CHILDREN

WARNING - EYE IRRITANT

CAUTION



POISON

Net Contents: 10 Litres

Gharda USA, Inc.

660 Newtown Yardley Road, Suite 106
Newtown, PA 18940
(215) 968-9474

Oracle® Dicamba **Agrcultural** Herbicide controls broadleaf weeds in cereals, corn, reduced tillage (prior to seeding and reduced tillage fallow), pastures and rangeland grasses, crop-free land (summer fallow and stubble), red fescue, canary grass, seedling grasses grown for seed and forage and low bush blueberries. Also for control of deciduous and coniferous brush species and broadleaf weeds in non-cropland areas such as roadsides, hydro, pipeline and railway rights-of-way, airports, military bases, turf, wasteland and similar non crop land areas.

GENERAL PRECAUTIONS:

1. Oracle Dicamba **Agrcultural** Herbicide should not be applied on or near desirable trees or plants.
2. Apply Oracle Dicamba **Agrcultural** Herbicide when air temperature is between 10 - 25°C. Do not apply when there is a risk of severe fall in night temperature after use.
3. Do not contaminate domestic or irrigation water. Thoroughly clean application equipment.
4. Do not treat areas where movement of the chemical into the soil or surface washing may bring Oracle Dicamba **Agrcultural** Herbicide into contact with roots of desirable plants.
5. Crop damage can occur if the chemical is applied at any time other than the recommended crop stage. (NOTE: Crops growing under stress from adverse environmental conditions such as excess moisture, drought, disease, etc., may suffer a further setback and exhibit more pronounced injury symptoms if Oracle Dicamba **Agrcultural** Herbicide is applied. However, the crop injury that may occur is usually offset by the weed control obtained).
6. Do not use additives such as oil, wetting agents, emulsifiers, detergents, spreaders, sticking agents or dispersing agents with Oracle Dicamba **Agrcultural** Herbicide on crops.
7. For information on feeding and grazing of beef and dairy cattle on treated vegetation and for recommendations on treatment/harvest intervals, refer to the table on Grazing Restrictions.
8. If Oracle Dicamba **Agrcultural** Herbicide is tank mixed with another product, such as 2,4-D, consult that product's label for additional safety precautions, restrictions, application rates, timings and additional weeds controlled.
9. Ensure that spray equipment used to apply Oracle Dicamba **Agrcultural** Herbicide is properly cleaned before reusing to apply any other chemicals. See section on suggested procedure for cleaning spray equipment.
10. **If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca**

SPRAY DRIFT PRECAUTIONS:

Oracle Dicamba **Agrcultural** Herbicide may cause injury to desirable trees and plants, particularly soybeans, flowers, fruit trees, grapes, ornamentals, peas, potatoes, tomatoes, tobacco and other broadleaf plants especially in their developmental and growing stage.

Follow these precautions when spraying in the vicinity of sensitive crops:

1. Avoid spraying when winds are gusty or in excess of 8 km/h and moving towards sensitive crops. Leave an adequate buffer zone between areas to be treated and sensitive plants.
2. Use coarse sprays since they are less likely to drift than fine sprays. Select nozzles which minimize amounts of the fine spray particles. Keep the spray pressure below 150 kPa and the spray volume above 220 L/ha unless otherwise required by the nozzle manufacturer.
3. Do not spray when the temperature is expected to exceed 30°C.
4. Avoid spraying under conditions of high humidity or fog.

RESISTANCE-MANAGEMENT RECOMMENDATIONS

For resistance management, Oracle Dicamba Agrcultural Herbicide is a Group 4 herbicide. Any weed population may contain or develop plants naturally resistant to Oracle Dicamba Agrcultural Herbicide and other Group 4 herbicides. The resistant biotypes may dominate the weed population if these herbicides are used repeatedly in the same field. Other resistance mechanisms that are not linked to site of action, but specific for individual chemicals, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

To delay herbicide resistance:

- Where possible, rotate the use of Oracle Dicamba Agrcultural Herbicide or other Group 4 herbicides with different herbicide groups that control the same weeds in a field.
- Use tank-mixtures with herbicides from a different group when such use is permitted.
- Herbicide use should be based on an IPM program that includes scouting, historical information related to herbicide use and crop rotation, and considers tillage (or other mechanical) , cultural, biological and other chemical control practices.
- Monitor treated weed populations for resistance development.
- Prevent movement of resistant weed seeds to other fields by cleaning harvesting and tillage equipment and planting clean seed.
- Contact your local extension specialist or certified crop advisors for any additional pesticide resistance-management and/or integrated weed-management recommendations for specific crops and weed biotypes.
- For further information or to report suspected resistance, contact Gharda USA, Inc. at 215-497-9501.

PRECAUTIONS:

1. KEEP OUT OF REACH OF CHILDREN.
2. Harmful if swallowed or absorbed through the skin.
3. Avoid contact with skin, eyes and clothing.
4. Thaw if frozen. Shake before use.

FIRST AID (Substituted Benzoic acid)	
If Inhaled:	<ul style="list-style-type: none"> ▪ Move person to fresh air. ▪ If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. ▪ Call a poison control centre or doctor for further treatment advice.
If on Skin or Clothing:	<ul style="list-style-type: none"> ▪ Take off contaminated clothing. ▪ Rinse skin immediately with plenty of water for 15-20 minutes ▪ Call a poison control centre or doctor for treatment advice.
If in Eyes:	<ul style="list-style-type: none"> ▪ Hold eye open and rinse slowly and gently with water for 15-20 minutes. ▪ Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. ▪ Call a poison control centre or doctor for further treatment advice.
If Swallowed:	<ul style="list-style-type: none"> ▪ Call a poison control center or doctor immediately for treatment advice. ▪ Have person sip a glass of water if able to swallow. ▪ Do not induce vomiting unless told to do so by a poison control centre or doctor. ▪ Do not give anything by mouth to an unconscious person.
HOT LINE NUMBER (Substituted Benzoic acid)	
Have the product container or label with you when calling a poison control centre or doctor, or going for treatment. For emergency medical treatment information call: 1-866-359-5660	
NOTE TO PHYSICIAN Probable mucosal damage may contraindicate the use of gastric lavage.	

TOXICOLOGICAL INFORMATION:

Treatment is symptomatic. Although symptoms are nonspecific, they may include exhaustion, muscular spasms, urinary incontinence, dyspnea and cyanosis. For ingestion, lavage stomach with tap water. Instill 30 gm activated charcoal in 90-120 ml. (3-4 oz.) of water. Catharsis with 15 gm sulfate in 175-240 ml. (6-8 oz.) water.

DIRECTIONS FOR USE

CEREALS

(not underseeded to legumes)

Treatment notes:

1. For best performance, spray when weeds are in the 2 to 3 leaf stage and rosettes are less than 5 cm across.
2. Use the higher level of listed rate ranges when treating more mature weeds or dense vegetative growth.
3. Crop damage can occur if application is made at any time other than the recommended crop stage.
4. Do not apply Oracle Dicamba **Agrcultural** Herbicide or Oracle Dicamba **Agrcultural** Herbicide tank mixes if crop is under-seeded to legumes.

Ground application:

Apply Oracle Dicamba **Agrcultural** Herbicide or Oracle Dicamba **Agrcultural** Herbicide tank mixes in at least 110 litres of water/ha.

Aerial application (Western Canada only):

Apply only by fixed-wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for the specific use, this product cannot be applied by any type of aerial equipment.

Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Use Precautions

Apply only when meteorological conditions at the treatment site allow for complete and even crop coverage. Apply only under conditions of good practice specific to aerial application as outlined in the *Basic Knowledge Requirements for Pesticide Education in Canada: Applicator Core and Aerial Module*, developed by CAPCO.

Do not apply to any body of water. Avoid drifting of spray onto any body of water or other **non-target** areas. Specified buffer zones should be observed.

Coarse sprays are less likely to drift, therefore, avoid combinations of pressure and nozzle type that will result in fine particles (mist). Do not apply during periods of dead calm or when wind velocity and direction pose a risk of spray drift. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, terrestrial habitat (such as shelter-belt) or aquatic habitat.

Operator Precautions

Do not allow the pilot to mix chemicals to be loaded onto aircraft. Loading of premixed chemicals with a closed system is permitted.

It is desirable that the pilot have communication capabilities at each treatment site at the time of application.

The field crew and the mixer/loaders must wear chemical resistant gloves, coveralls and goggles or face shield during mixing/loading, cleanup and repair. Follow the more stringent label precautions in cases where the operator precautions exceed the generic label recommendations on the existing ground boom label.

All personnel on the job site must wash hands and face thoroughly before eating and drinking. Protective clothing, aircraft cockpit and vehicle cabs must be decontaminated regularly.

Product Specific Precautions

Read and understand the entire label before opening this product. If you have questions, call the manufacturer at 1-215-968-9474 or obtain technical advice from the distributor or your provincial agricultural representative. Application of this specific product must meet and/or conform to the following:

1. Oracle Dicamba Agricultural Herbicide or Oracle Dicamba Agricultural Herbicide phenoxy herbicide tank mixes may be aurally applied in not less than 20 litres of water/ha.
2. Apply Oracle Dicamba Agricultural Herbicide alone at 230 mL/ha or tank mix Oracle Dicamba Agricultural Herbicide at 230 mL/ha with the recommended rate of the phenoxy herbicides specified on this label.
3. Do not spray when wind velocities are greater than 15 km/hour.
4. Do not use nozzle pressure above 200 kPa.
5. Do not spray when the wind is blowing towards a nearby sensitive crop, garden, or shelterbelt.

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agricultural Herbicide Rate	Tank Mix
buckwheat, Tartary, buckwheat, wild cockle, cow cleavers (higher rate only) lady's thumb, sow-thistle, perennial (top growth only) smartweed, green spurry, corn thistle, Canada (top growth only)	Oracle Dicamba Agricultural Herbicide alone at 230-290mL/ha	none
All of the above plus: burdock (young seedlings) canola, volunteer* cocklebur flixweed hemp-nettle** kochia pigweed, <i>redroot</i> pigweed, <i>Russian</i> radish, wild shepherd's-purse sunflower, <i>volunteer</i> *** thistle, <i>Russian</i>	Oracle Dicamba Agricultural Herbicide at 230 mL/ha +	2,4-D amine or MCPA amine or MCPA K
All of the above plus: chickweed hemp-nettle** spurry, corn; stinkweed; sunflower, <i>volunteer</i> ***	Oracle Dicamba Agricultural Herbicide at 230 mL/ha +	Sencor or Lexone
All of the above plus: buckwheat, wild; canola, volunteer;* sow-thistle, perennial (top growth only); thistle, Canada (top growth only)	Oracle Dicamba Agricultural Herbicide at 230 mL/ha +	Ally

* Best results will be obtained if application is made prior to bolting of canola, when this weed is at the 2 to 4 leaf stage.

** Use Oracle Dicamba Herbicide + MCPA K for hemp-nettle control. Apply at the 2 to 3 leaf stage of weed for best control. Hemp-nettle may not be controlled if application is made at a more advanced stage of crops and weeds.

*** Depending on the growing conditions, control may be slightly delayed.

APPLICATION DIRECTIONS FOR CEREALS:

Oracle Dicamba Agricultural Herbicide may be applied to:

- Spring Wheat
- Spring Barley
- Winter Wheat
- Oats
- Spring Rye

The following sections describe application directions for these crops.

Spring wheat:

Herbicide Mix	Rate/ha	Crop Stage
Oracle Dicamba Agricultural Herbicide alone	230-290 mL/ha	2 - 5 leaf
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	2 - 5 leaf
or MCPA amine	850 mL/ha (500 g/L formulation)	2 - 5 leaf
or MCPA K	1.1 L/ha (400 g/L formulation)	2 - 5 leaf
or Sencor ® 500*	275-425 mL/ha**	2 - 3 leaf
or Lexone ® DF*	275 g/ha	2 - 3 leaf
or Ally***	5 g/ha	2 - 5 leaf

* Sencor/Lexone tank mixes apply to Western Canada only. Application may be delayed until the 4-leaf stage of the crop, however, crop tolerance may be reduced. Apply Oracle Dicamba Agricultural Herbicide at 230 mL/ha with Sencor/Lexone.

** Use the higher rate of Sencor 500 for control of volunteer sunflowers.

*** Ally tank mixes apply to Western Canada only. Apply Oracle Dicamba Agricultural Herbicide at 230 mL/ha with Ally. Ensure that Ally ® is completely in suspension in the spray tank before adding Oracle Dicamba Herbicide. Do not add a surfactant.

Spring rye:

Herbicide Mix	Rate/ha	Crop Stage
Oracle Dicamba Agricultural Herbicide	230-290 mL/ha	2 - 3 leaf
alone		
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	2 - 3 leaf

Spring barley:

Herbicide Mix	Rate/ha	Crop Stage
Oracle Dicamba Agricultural Herbicide alone	230-290 mL/ha	2 - 5 leaf
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	2 - 5 leaf
or MCPA amine	850 mL/ha (500 g/L formulation)	2 - 5 leaf
or MCPA K	1.1 L/ha (400 g/L formulation)	2 - 5 leaf
or Sencor 500*	275-425 mL/ha**	2 - 3 leaf
or Lexone DF*	275 g/ha	2 - 3 leaf
or Ally***	5 g/ha	2 - 5 leaf

* Sencor/Lexone tank mixes apply to Western Canada only. NOTE: Do not use on Klondike barley.

** Use the higher rate of Sencor 500 for control of volunteer sunflowers.

*** Ally tank mixes apply to Western Canada only. Apply Oracle Dicamba Herbicide at 230 mL/ha with Ally. Ensure that Ally ® is completely in suspension in the spray tank before adding Oracle Dicamba Agricultural Herbicide.

Winter wheat:

Herbicide Mix	Rate/ha	Crop Stage
Oracle Dicamba Agricultural Herbicide alone	230-290 mL/ha	15 - 25 cm tall or before shot blade stage
+ 2,4-D amine	850 mL/ha (500 g/L formulation)	15 - 25 cm tall or before shot blade stage
or MCPA amine	850 mL/ha (500 g/L formulation)	
or MCPA K	1.1 L/ha (400 g/L formulation)	

Oats:

Herbicide Mix	Rate/ha	Crop Stage
Oracle Dicaamba Agricultural Herbicide alone	230-290 mL/ha	2 - 5 leaf
+ MCPA amine	850 mL/ha (500 g/L formulation)	2 - 5 leaf
or MCPA K	1.1 L/ha (400 g/L formulation)	2 - 5 leaf

GRAZING RESTRICTIONS:

1. Following treatment with Oracle Dicamba Agricultural Herbicide plus any other herbicide tank mix: Do not graze or harvest for livestock feed prior to crop maturity; sufficient data are not available to support such use.
2. Following treatment with Oracle Dicamba Agricultural Herbicide or Oracle Dicamba Agricultural Herbicide plus 2,4-D, follow the grazing restrictions below:

Hay making and grazing restrictions for dairy cattle:

Rate/ha	Delay between treatment and hay-making/grazing
Up to 1.25 L Oracle Dicamba Agricultural Herbicide	9 days

Hay making, grazing and slaughter restrictions for meat animals:

Number of days between Oracle Dicamba Agricultural Herbicide application and hay-making or consumption	Slaughter restrictions
Vegetation consumed within 30 days of treatment	feed animal untreated diet for 30 days before slaughter
Vegetation consumed 30 days after treatment	no restrictions on slaughter

FIELD CORN:**Treatment Notes**

1. Apply Oracle Dicamba Agricultural Herbicide or Oracle Dicamba Agricultural Herbicide tank mixes in 220 litres of water/ha at a pressure of 150 to 275 kPa. Use coarse sprays.
2. Keep spray mixture in suspension at all times. If mixture is allowed to settle, thoroughly agitate the mixture before spraying.
3. Do not apply to sweet corn.
4. Do not use additives. Oils, wetting agents, emulsifiers, detergents, spreaders, sticking agents, or dispersing agents are not recommended with Oracle Dicamba Agricultural Herbicide.
5. Corn height refers to the crop as it stands, not leaf-extended.
6. When using drop pipes (drop nozzles), direct the spray beneath the lower leaves of the corn and onto the weeds and soil. Do not apply to corn over 50 cm in height.
7. Apply no later than 2 weeks prior to tassel emergence when using Oracle Dicamba Agricultural Herbicide alone up to 50 cm.
8. For the best control of annuals, spray when they are actively growing and in the seedling stage. Poor results may occur if weeds are well advanced at the time of application.
9. When applying Oracle Dicamba Agricultural Herbicide adjacent to sensitive crops, apply as a pre-emergent or early post-emergent treatment to avoid potential drift onto these sensitive crops.
10. If this pest control product is to be used on a commodity that may be exported to the U.S. and you require information on acceptable residue levels in the U.S., visit CropLife Canada's web site at: www.croplife.ca

DO NOT APPLY BY AIR.

ORACLE DICAMBA HERBICIDE / LIQUID NITROGEN:

Pre-emergent applications of Oracle Dicamba Agricultural Herbicide are generally compatible with most liquid nitrogen fertilizers. To determine compatibility, mix all components of the finished spray in proportionate quantities in a small jar before mixing in the spray tank. If the herbicides do not ball-up or form flakes, sludge, jelly, oily films or layers, or other precipitates within 5 minutes after mixing, the tested spray-mix is compatible.

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agricultural Herbicide Rate	Tank Mix
bindweed, field** buckwheat, Tartary buckwheat, wild cleavers cockle, cow lady's-thumb, lamb's-quarters* mustard, hare's-ear mustard, Indian mustard, tumble mustard, wild mustard, wormseed pigweed, redroot* pigweed, Russian ragweed, common* ragweed, false ragweed, giant sow-thistle, perennial** spurry, corn smartweed, green thistle, Canada** velvetleaf	Oracle Dicamba Herbicide alone at 600 mL - 1.25 L/ha	none

* Including atrazine-resistant species.

** Apply Oracle Dicamba Agricultural Herbicide annually for three years at the flowering stage of bindweed and the budding stage of thistles.

PRE-EMERGENCE TREATMENT

EASTERN CANADA ONLY

Oracle Dicamba **Agrcultural** Herbicide can be used alone at 1.25 L/ha or in tank mixes with the following herbicides for additional broadleaf and grassy weed control.

Herbicide	Rate/ha
Dual	2.0 - 2.75 L
Dual II	2.0 - 2.75 L
Frontier	1.1 - 1.4 L
Primextra Light	5.80 - 7.70 L
Atrazine 480*	2.10 L
Prowl 400**	4.20 L
Atrazine 480* + Dual II	2.10 L + 2.0 L

* Other atrazine formulations will require a rate calculation adjustment according to percent active ingredient.

** Other pendimethalin formulations will require a rate calculation adjustment according to percent active ingredient.

Pre-emergence treatment notes:

- Apply Oracle Dicamba **Agrcultural** Herbicide tank mixes as broadcast ground treatments after planting but before weeds and corn emerge.
- Apply to medium to fine textured soils containing more than 2.5% organic matter.
- Do not use on sandy or sandy loam soils.
- Avoid direct chemical contact with the corn seed. If you plan to apply Oracle Dicamba **Agrcultural** Herbicide prior to corn emergence, be sure to place the corn seeds 4 cm or more below the soil surface. If seeds are planted less than 4 cm below the soil surface, delay application of Oracle Dicamba Herbicide until the spike stage.
- Do not incorporate. If applications are made during planting, apply Oracle Dicamba **Agrcultural** Herbicide far enough behind the planting equipment to avoid incorporation by the planter wheel or other covering device. If soil crusting makes it necessary to use a rotary hoe after a pre-emergence treatment, delay hoeing the soil more than 1.3 cm deep.
- Always consult the tank mix partner label for further limitations and restrictions (especially re: soil type).

POST-EMERGENCE TREATMENT:

Oracle Dicamba Agrcultural Herbicide or Oracle Dicamba Agrcultural Herbicide tank mixes can be applied as “overlays” to corn previously treated with any other broadleaf or grass herbicide. The 1.25 L rate of Oracle Dicamba Agrcultural Herbicide as “overlay” is particularly effective in controlling velvetleaf and providing extended residual control of other late germinating, deep rooted annuals.

Note: Do not use additives such as oils, wetting agents, or sticking agents.

Oracle Dicamba Herbicide alone, Spike to 5-leaf corn, Eastern and Western Canada			
Herbicide	Rate/ha	Corn Stage	Weed Stage
Oracle Dicamba Agrcultural Herbicide alone	1.25 L/ha	Emergence to 5-leaf	Pre-emergence to 2-leaf (1)

- (1) For best performance, spray when the broadleaf weeds are emerged and up to the 2-leaf stage of their development.

Oracle Dicamba Herbicide tank mixes:**Eastern Canada only**

Herbicide	Rate/ha	Corn Stage	Weed Stage
Oracle Dicamba Agrcultural Herbicide + Frontier®	1.25 L/ha + 1.1 - 1.4 L	Spike to 3-leaf	Pre-emergence to 2-leaf ***
Oracle Dicamba Agrcultural Herbicide + Atrazine 480*	1.25 L + 2.10 L	Spike to 5-leaf	Pre-emergence to 2-leaf
Oracle Dicamba Agrcultural Herbicide + Atrazine 480* + Dual II	0.6 - 1.25 L + 2.3 L + 2.0 - 2.75 L	Spike to 2-leaf	Emergence to 2-leaf
Oracle Dicamba Agrcultural Herbicide + Primextra® Light	0.6 - 1.25 L + 5.80 + 7.70 L	Spike to 2-leaf	Emergence to 2-leaf
Oracle Dicamba Agrcultural Herbicide + Prowl® 400**	0.6 - 1.25 L + 4.20 L	Spike to 4-leaf	Pre-emergence to 2-leaf
Oracle Dicamba Agrcultural Herbicide + Ultim® 37.4 DF + non-ionic surfactant	0.6 L + 1 bag + 0.2% v/v	Spike to 6-leaf	Emergence to 6-leaf
Oracle Dicamba Agrcultural Herbicide + Elim® EP + non-ionic surfactant	0.6 L + 60g + 0.2% v/v	Spike to 3-leaf	Emergence to 4-leaf
Oracle Dicamba + Dual II	0.6 - 1.25 L + 2.0 - 2.75 L	Spike to 2-leaf	Emergence to 2-leaf
Oracle Dicamba Agrcultural Herbicide + Prowl® 400** + Elim EP + non-ionic surfactant	0.625 L + 2.5 L + 50 g + 0.2% v/v	Spike to 3-leaf	Emergence to 4-leaf

* Other atrazine formulations will require a rate calculation adjustment according to percent active ingredient.

** Other pendimethalin formulations will require a rate calculation adjustment according to percent active ingredient.

*** For annuals, apply before 2-leaf stage.

**Spike to 50 cm standing corn
Eastern and Western Canada only**

Herbicide	Rate/ha	Corn Stage	Weed Stage
Oracle Dicamba Agrcultural Herbicide alone	600 mL	Emergence to 50 cm (drop to nozzles from 20-50 cm corn)	Pre-emergence to 2- leaf
Oracle Dicamba Agrcultural Herbicide + 2,4-D amine	290 mL + 850 mL	Emergence to 50 cm (drop nozzles from 20-50 cm corn)	Pre-emergence to 2- leaf

**Sequential Oracle Dicamba Agrcultural Herbicide applications
Eastern and Western Canada**

Oracle Dicamba Agrcultural Herbicide may be applied sequentially to a Oracle Dicamba Agrcultural Herbicide application to control late-emerging weeds such as field bindweed, Canada thistle and velvetleaf. Follow application directions as outlined for the Oracle Dicamba Herbicide alone post-emergence treatments up to 50 cm tall corn.

Grazing restrictions

Do not graze cattle on treated crop, or harvest for silage until 7 days following the application of Oracle Dicamba Agrcultural Herbicide alone and at least 12 weeks following treatment with Oracle Dicamba Agrcultural Herbicide tank mixes.

WEED CONTROL IN REDUCED TILLAGE (PRIOR TO SEEDING)

Treatment notes:

1. Oracle Dicamba Agrcultural Herbicide + Roundup® applications may be applied to emerged annual grass and annual broadleaf weeds in reduced tillage systems prior to seeding of wheat, barley, rye, oats and field corn only.
2. Do not apply prior to seeding sweet corn.
3. Planting should follow soon after application since this tank mix does not provide residual weed control.
4. Delayed planting following chemical application will allow weeds to emerge between application and crop emergence.
5. For field corn, apply to medium to fine textured soils containing more than 2.5% organic matter. Do not use on sandy or sandy loam soils.
6. Certain broadleaf crops such as sweet corn, lentils, peas, canola and flax can be injured by a pre-seeding application of this tank mix and should not be planted after the use of this tank mix.
7. Under certain stress conditions, such as drought, cool temperatures where extremely hard water (>700ppm CA+ Mg) will be used, use 50L/ha of water with this tank mix to help improve results.

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agrcultural Herbicide	Tank Mix
Annual grasses (Apply any time between emergence and heading) brome, downy cereals, volunteer darnel, Persian foxtail, green oats, wild	Rate Oracle Dicamba Agrcultural Herbicide at 315 mL/ha +	Roundup® at 935 mL/ha + 0.5 L of a non-ionic surfactant in 100 L of water
Annual broadleaves (Apply up to 15 cm height) buckwheat, wild* canola, volunteer cockle, cow flixweed** kochia lady's-thumb lamb's-quarters mustard, wild pigweed, redroot smartweed stinkweed** thistle, Russian	Oracle Dicamba Agrcultural Herbicide at 315 mL/ha +	Roundup® at 935 mL/ha + 0.5 L of a non-ionic surfactant in 100 L of water
Perennials (Apply before initiation of seed head or browning of lower leaves) barley, foxtail (suppression only)	Oracle Dicamba Agrcultural Herbicide at 315 mL/ha +	Roundup® at 935 mL/ha + 0.5 L of a non-ionic surfactant in 100 L of water

* Apply at the 1 to 4-leaf stage.

** For optimal control of winter annual broadleaf weeds such as flixweed and stinkweed, 2,4-D should be applied to emerged, actively growing weeds in the fall the year prior to the Oracle Dicamba Agrcultural Herbicide + Roundup spring pre-seeding tank mix. Refer to the 2,4-D product label for appropriate rates.

WEED CONTROL IN REDUCED TILLAGE FALLOW

Treatment notes:

1. Apply Oracle Dicamba **Agricultural** Herbicide tank mixes in the spring to fallow land when seedling weeds have emerged, and are actively growing at the 2 to 4-leaf stage.
2. Reduced control may occur if applications are made at an advanced stage of weed development.

DO NOT APPLY BY AIR.

Weeds Controlled (in alphabetical order)	Rate	Oracle Dicamba Agricultural Herbicide Tank Mix
buckwheat, wild buckwheat, Tartary cockle, cow flixweed kochia lady's-thumb lamb's-quarters mustard, wild pigweed, redroot shepherd's-purse smartweed, green sow-thistle perennial (top growth) thistle, Canada (top growth) thistle, Russian	230 - 290 mL/ha +	1.1 L/ha of 2,4-D amine 500 OR 920 mL/ha of 2,4-D L.V. ester 600 in 50-100 L of water
barley, foxtail** buckwheat, wild** cereals, volunteer cockle, cow flixweed* foxtail, green kochia lady's-thumb lamb's-quarters mustard, wild oats, wild pigweed, redroot** canola, volunteer stinkweed thistle, Russian	290 mL/ha +	750 mL -1.0 L/ha Roundup® + 350 mL of non-ionic surfactant registered for use in 50 - 100L of water
buckwheat, wild	600 mL/ha +	750 mL -1.0 L/ha Roundup® + 350mL of non-ionic surfactant registered for use in 50 – 100 L of water

* For control of flixweed use 1.0 L/ha of Roundup®

** Suppression only.

Oracle Dicamba Agrcultural Herbicide/Roundup® application notes:

1. These tank mixes should be applied to emerged actively growing annual weeds from 8-15 cm in height.
2. Use the higher rate of Roundup when weeds are at a more advanced stage of growth.
3. For perennial weed control, refer to the appropriate section of this label for proper stages of growth and recommended stages of application.
4. Reduced control may occur if muddy water is used, such as water from dug-outs, ponds and unlined ditches.

PERENNIAL WEED CONTROL IN SUMMERFALLOW AND STUBBLETreatment notes:

1. Apply Oracle Dicamba Agrcultural Herbicide in 110-220 litres of water/ha.
2. For the most effective control of Canada thistle, follow a long-term approach that includes in-crop, post-harvest, and summer fallow treatments, in conjunction with tillage operations.
3. If application is made after September 1st, or if soil moisture levels are extremely low after application, crop injury may occur in the spring following application.

DO NOT APPLY BY AIR.

Weeds Controlled (in alphabetical order)	Rate	Recropping in year following
bindweed, field daisy, English dock, curled (top growth) goldenrod ragwort, tansy sow thistle, perennial thistle, Canada	Oracle Dicamba Agrcultural Herbicide alone at 2.5 L/ha	cereals soybeans field corn white beans sweet corn
thistle, Canada sow-thistle, perennial	Oracle Dicamba Agrcultural Herbicide at 1.25 L/ha + Roundup® at 1.7 L/ha +	All of the above plus canola

350 mL of a non-ionic surfactant per 100 L of water

Application directions

Summerfallow treatment notes:

1. Cultivate in the spring and apply Oracle Dicamba **Agricultural** Herbicide when:

Weed (in alphabetical order)	Weed Stage
thistles	the majority of thistles are up and before the early bud stage (15-25 cm tall)
field bindweed	in the flowering stage
other weeds	in the early bud stage of growth

2. Cultivate three weeks after application.

Stubble treatment notes:

1. Apply to regrowth after harvest and at least 2 weeks prior to a killing frost.

Perennial rosette control in summerfallow

Treatment notes:

1. For the most effective control of Canada thistle, follow a long-term approach that includes in-crop, post-harvest, and summerfallow treatments, in conjunction with tillage operations.
2. Commence early spring cultivation and continue as required throughout the summer. Note: The final cultivation must occur by the end of July between July 15 - August 1 and the final cultivation should cut the thistle off 5 to 7.5 cm below the soil surface.
3. Spray in 110-220 L of water/ha when the majority of thistles have emerged as low growing rosettes 15 to 25 cm across.
4. Apply at least two weeks prior to a killing frost.
5. Cultivate three weeks after application.

DO NOT APPLY BY AIR.

Weeds Controlled	Rate	Recropping in year following
thistle, Canada	1.25 L/ha	cereals field corn white beans canola soybeans

PASTURES, RANGELAND, AND NON-CROP AREAS

Treatment notes:

For broadleaf weed control:

1. Apply Oracle Dicamba **Agrcultural** Herbicide or Oracle Dicamba **Agrcultural** Herbicide tank mixes in 1 of water/ha when weeds are actively growing. Thorough coverage of foliage is necessary to control weeds.
2. Do not apply Oracle Dicamba **Agrcultural** Herbicide or Oracle Dicamba **Agrcultural** Herbicide tank mix if pasture is underseeded to legumes.

DO NOT APPLY BY AIR.

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agrcultural Herbicide	Tank mix
bindweed, field daisy, English dock, curled (top growth) goldenrod ragwort, tansy sow-thistle, perennial thistle, Canada	Rate Oracle Dicamba Agrcultural Herbicide alone at 2.1 L/ha	none
beard, goat's cherry, ground knapweed, diffuse sage, pasture sorrel, sheep spurge, thyme-leafed weed, poverty	Oracle Dicamba Agrcultural Herbicide alone at 4.6 L/ha	none
poison ivy	Oracle Dicamba Agrcultural Herbicide at 1.65 L/ha +	2.2 L/ha of 2,4-D amine (500 g/L formulation) in 560 L of water/ha
wild carrot plus additional weeds found on the 2,4-D amine label	Oracle Dicamba Agrcultural Herbicide 2.1 L/ha +	2.2 L/ha of 2,4-D amine (500 g/L formulation)
All of the above plus: additional weeds found on the 2,4-D amine label	Oracle Dicamba Agrcultural Herbicide 2.1 L/ha +	1.83 L/ha of 2,4-D L.V. ester (600 g/L formulation)

For brush weed control:

1. Oracle Dicamba **Agrcultural** Herbicide is effective in controlling many deciduous brush species that are found growing along fence rows and in other areas around the farm where they may be undesirable.
2. Apply Oracle Dicamba **Agrcultural** Herbicide tank mixes in spring or early summer to deciduous species (leaves should be fully expanded) either as a leaf stem treatment or as a broadcast ground application.
3. Brush and trees over 2 meters tall should be cut and regrowth treated when it develops.
4. Do not apply Oracle Dicamba **Agrcultural** Herbicide tank mixes if pasture or rangeland is underseeded to legumes.
5. For Stem Foliage Treatment, apply to all foliage and stems to the point of runoff. The volume of spray mix applied per hectare will vary according to the height and density of the woody species present.
6. For Broadcast Ground Treatment, apply Oracle Dicamba **Agrcultural** Herbicide tank mixes in sufficient dilution to wet all foliage. Normally, 220-230 litres of water/ha is recommended for brush stands.

DO NOT APPLY BY AIR.

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agrcultural Herbicide	Rate	Tank Mix
alder aspen poplar cherry western snowberry (buckbrush) wolf willow (silverwillow)	Oracle at 2.1 L/ha/1000 L of water +	4.0 L/ha of 2,4-D amine (500 g/L formulation) OR 3.3 L/ha of 2,4-D L.V. (600 g/L formulation)	
aspens poplar	Oracle Dicamba Agrcultural Herbicide at L/ha +	3.4 L/ha of 2,4-D amine (500 g/L formulation) OR 3.75 L/ha of 2,4-D L.V. ester (600 g/L formulation)	
prickly rose	Oracle Dicamba Agrcultural Herbicide at L/ha +	3.4 L/ha of 2,4-D amine (500 g/L formulation) OR 3.75 L/ha of 2,4-D L.V. ester (600 g/L formulation)	
western snowberry	Oracle Dicamba Agrcultural Herbicide at L/ha +	3.3 L/ha of 2,4-D L.V. ester (600 g/L formulation)	

Grazing restrictions:

Hay making and grazing restrictions for dairy cattle:

Rate/ha	Delay between treatment and hay making/grazing
Up to 1.25 L Oracle Dicamba Agrcultural Herbicide	0 days
> 1.25 L to 2.3 L Oracle Dicamba Agrcultural Herbicide	7 days
> 2.3 L to 4.6 L Oracle Dicamba Agrcultural Herbicide	14 days
> 4.6 L to 7.1 L Oracle Dicamba Agrcultural Herbicide	30 days

Hay making, grazing and slaughter restrictions for restrictions for meat animals:

Number of days between Oracle Dicamba Agrcultural Herbicide application and hay making or consumption	Slaughter Restrictions
vegetation consumed within 30 days of treatment	feed animal untreated diet for 30 days before slaughter
vegetation consumed 30 days after treatment	no restrictions on slaughter

SEED PRODUCTION:

Treatment notes:

For new/established stands of red fescue

1. Apply Oracle Dicamba Agrcultural Herbicide or Oracle Dicamba Agrcultural Herbicide tank mixes in at least 110 litres of water/ha.
2. Applications to new seedling stands may be made when the crop is 5 cm tall.
3. Application to established stands may be made up to the shot-blade stage of the crop.
4. For dandelion control, apply Oracle Dicamba Agrcultural Herbicide plus 2,4-D amine in the fall when weeds are in the rosette or early bud stage.

DO NOT APPLY BY AIR.

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agrcultural Herbicide Rate	Tank Mix
buckwheat, wild buckwheat, Tartary cockle, cow clover lady's-thumb sow-thistle, perennial (top growth) spurry, corn smartweed, green thistle, Canada (top growth)	Oracle Dicamba Agrcultural Herbicide alone at 600 mL/ha	none
All of the above plus: additional weeds found on the 2,4-D amine label	Oracle Dicamba Agrcultural Herbicide at 600 mL/ha +	1.5 L/ha of 2,4-D amine (500 g/L formulation)

For canary grass:

1. The canary grass should only be used as bird seed.
2. For specific weeds controlled, refer to the Oracle Dicamba Agrcultural Herbicide + MCPA amine weed spectrum list under "Cereals"

Herbicide	Rate	Canary Grass Stage
Oracle Dicamba Agrcultural Herbicide		
Oracle Dicamba Agrcultural Herbicide + MCPA amine (500 g/L formulation)	290 mL/ha 290 mL/ha + 850 mL/ha	3 - 5 leaf stage 3 - 5 leaf stage

For seedling grasses (seeded alone or underseeded with cereals):

For seed and forage production of the following seedling grasses:

bromegrass, smooth	wheatgrass, crested
fescue, meadow	wheatgrass, intermediate
fescue, tall	wheatgrass, pubescent
foxtail, meadow	wheatgrass, slender
orchard grass	wheatgrass, streambank
red fescue, creeping	wheatgrass, tall
timothy	

1. Apply Oracle Dicamba Herbicide or Oracle Dicamba Agrcultural Herbicide + tank mixes in at least 110 litres of water/ha.
2. Application to new seedling grasses may be made when they are in the 2 to 4-leaf stage. If the seedling grass is under seeded with a cereal crop, refer to "Cereals" for additional restrictions pertaining to application type and rate.
3. If the crops are to be used as feed or pasture following treatment with Oracle Dicamba Agrcultural Herbicide, Oracle Dicamba Agrcultural Herbicide plus 2,4-D amine or MCPA, refer to "Grazing Restrictions".

Weeds Controlled (in alphabetical order)	Oracle Dicamba Agrcultural Herbicide Rate	Tank Mix
buckwheat, Tartary buckwheat, wild cockle, cow cleavers (higher rate only) lady's-thumb sow-thistle, perennial (top growth) smartweed, green spurry, corn thistle, Canada (top growth)	Oracle Dicamba Agrcultural Herbicide alone at 230-290 mL/ha	none
All of the above plus: burdock (young seedlings) canola, volunteer* cocklebur flixweed hemp-nettle** kochia pigweed, redroot pigweed, Russian radish, wild shepherd's-purse sunflower, volunteer*** thistle, Russian	Oracle Dicamba Agrcultural Herbicide at 230-290 mL/ha	850 mL/ha of 2,4-D amine (500 g/L formulation) OR 850 mL/ha of MCPA amine (500 g/L formulation) OR 1.1 L/ha of MCPA K (400 g/L formulation)

* Best results will be obtained if application is made prior to bolting of canola, when this weed is at the 2 to 4-leaf stage.

** Use Oracle Dicamba Agrcultural Herbicide + MCPA K for hemp-nettle control. Apply at the 2 to 3-leaf stage of weed for best control. Hemp-nettle may not be controlled if application is made at a more advanced stage of crops and weeds.

*** Depending on the growing conditions, control may be delayed slightly.

For established grass pasture:

1. Apply Oracle Dicamba Agrcultural Herbicide at 600 mL/ha with 1.5 L/ha of 2,4-D amine (500 g/L formulation) to suppress volunteer alfalfa.
2. Apply Oracle Dicamba Agrcultural Herbicide + 2,4-D amine in 110 - 220 L/ha in the spring to actively growing alfalfa at greater than 5 cm in height.

LOW-BUSH BLUEBERRIES:Treatment notes:

1. Oracle Dicamba **Agrcultural** Herbicide can be used alone or in a tank mix with 2,4-D L.V. ester.
2. Apply Oracle Dicamba **Agrcultural** Herbicide or the Oracle Dicamba **Agrcultural** Herbicide tank mix in 550 litres of water per hectare.
3. Apply in the fall while the sweet-fern is still moderately green after 90% of the blueberries have dropped their leaves. This should be done before the area is burned. Fall burning or cutting should be carried out 4 to 5 weeks after spraying. If spring burning or cutting is planned, it should be done as early as possible in the spring to reduce injury to the blueberries.

DO NOT APPLY BY AIR.

Weeds Controlled	Oracle Dicamba Agrcultural Herbicide	Tank mix
	Rate	
fern, sweet	4.6 - 7.1 L/ha	none

lambkill (sheep laurel)

additional broadleaf control

2.3 L/ha +

no

5.7 L of 2,4-D L.V. ester
(600 g/L formulation)

NON CROP AREAS SUCH AS RIGHTS-OF-WAY, UTILITY, ROADSIDES, HYDRO, PIPELINE AND RAILWAY RIGHTS-OF-WAY, AIRPORTS, MILITARY BASES, TURF, WASTELAND AND SIMILAR NON CROP AREAS.

BRUSH CONTROL:

Oracle Dicamba Herbicide tank-mixtures are effective in controlling many coniferous and deciduous brush species found in areas where they may be undesirable.

1. Prepare a tank-mix consisting of Oracle Dicamba Agricultural Herbicide plus 2,4-D amine, 2,4 D L.V. ester or 2,4 D/2,4 DP herbicide.
2. Apply Oracle Dicamba Agricultural Herbicide tank-mixes in spring or early summer to coniferous and deciduous brush species (after the leaves have fully expanded). Stop applications at least 3 weeks prior to change of leaf colour in the fall.
3. For best results and ease of application, cut brush or trees over 2M in height and treat re-growth when it develops.
4. Do not mix Oracle Dicamba Agricultural Herbicide with any oils.

FOLIAGE AND STEM TREATMENT:

Ground Application Directions

1. Apply to foliage and to stems to the point of run-off. Complete coverage is essential for effective control.
2. The volume of spray required per hectare will vary according to the height and density of the woody species present.
3. Use of a tank-mix with 2,4 D L.V. ester herbicide with the water conditioner Adjust may improve the control of balsam poplar as well as other species when they are under drought stress, especially under periods of hot, dry weather.

SPECIES CONTROLLED	RATE PER 1000L OF SPRAY WATER
Wolf willow (silver willow) Aspen poplar Wild rose Alder, Cherry	2.1 L of Oracle Dicamba Agricultural Herbicide + 4L of 2,4-D amine(500g/L formulation) herbicide or 3.3 L of 2,4-D L.V. ester (600g/L formulation) herbicide
Balsam poplar Basswood Birch Elm Hickory Vine maple Black cottonwood Bur Oak Red Oak Tamarack Pine White Cedar Spruce Balsam fir	4 L of Oracle Dicamba Agricultural Herbicide + 8 L 2,4-D amine (500 g/L formulation) herbicide or 6.6 L of 2,4-D L.V. ester 600g/L formulation) herbicide
Sugar maple White ash	5.2 L Oracle Dicamba Agricultural Herbicide plus 7.1 L of 2,4-D/2,4- DP L.V. ester (350 g/L 2,4-D; 350 g/L 2,4-DP formulation)

BROADCAST GROUND TREATMENT:

Application Directions:

1. Apply in sufficient water to wet all foliage
2. A water volume of 220-330 Litres per hectare is recommended for bush stands.

SPECIES CONTROLLED	RATE PER HECTARE
White (paper) birch Willow Aspen poplar	4.2 L of Oracle Dicamba Agricultural Herbicide plus 8 L of 2,4-D amine (500g/L formulation) herbicide or 6.6 L 2,4-D L.V.ester (600g/L formulation) herbicide or 5.2 L of Oracle Dicamba Agricultural Herbicide + 7.2 L of 2,4-D/2,4-DP L.V. ester (350g/L 2,4-D; 350 g/L 2,4-DP formulation) herbicide

AERIAL APPLICATIONS:

Apply only by fixed wing or rotary aircraft equipment which has been functionally and operationally calibrated for the atmospheric conditions of the area and the application rates and conditions of this label.

Label rates, conditions and precautions are product specific. Read and understand the entire label before opening and using this product. Apply only at the rate recommended for aerial application on this label. Where no rate for aerial application appears for a specific use, this product cannot be applied by any type of aerial equipment. Ensure uniform application. To avoid streaked, uneven or overlapped application, use appropriate marking devices.

Application Directions:

1. Oracle Dicamba Agricultural Herbicide tank-mixes may be applied aerially in not less than 85 litres of water per hectare
2. Do not spray when wind velocities are greater than 15 km/hour
3. Do not use nozzle pressure above 300 kPa.
4. Do not spray when wind is blowing towards nearby desirable trees or plants.

SPECIES CONTROLLED	RATE PER HECTARE
Aspen poplar White (paper) birch	4.2 L Oracle Dicamba Agricultural Herbicide + 8 L of 2,4-D amine (500g/L formulation) or 6.6 L of 2,4-D L.V. ester (600g/L formulation) herbicide.

Refer to Aerial Application Use Precautions and Operator Precautions under the "Cereals" section.

BROADLEAF WEED CONTROL

Oracle Dicamba Agricultural Herbicide can be used alone or in tank-mixes with 2,4-D amine or 2,4-D L.V. ester herbicide. Oracle Dicamba Agricultural Herbicide tank-mixes offer additional broadleaf weed control.

IMPORTANT (Ground Application Directions)

Apply Oracle Dicamba Agricultural Herbicide or Oracle Dicamba Agricultural Herbicide tank-mixes in 110 to 220 litres of water per hectare when weeds are actively growing. For poison ivy control use 560 litres of water per hectare. Thorough coverage of foliage is necessary to control weeds. For control of a broader range of weeds, the recommended rate of Oracle Dicamba Agricultural Herbicide may be tank-mixed with 2,4-D amine (500g/L) or 2,4-D ester (600 g/L) herbicide. Consult 2,4-D herbicide label regarding rates of application and weed species controlled.

SPECIES CONTROLLED (alphabetical order)	RATE PER HECTARE
Poison Ivy	1.7 L Oracle Dicamba Agricultural Herbicide + 2.2 L of 2,4-D Amine (500g/L) herbicide
Wild Carrot	2.1 L Oracle Dicamba Agricultural Herbicide + 4.4 L of 2,4-D Amine (500g/L) herbicide
<i>Oracle Herbicide alone for top growth control of:</i> Absinthe Canada thistle Perennial Sow Thistle Leafy Spurge Poverty Weed Scentless mayweed	1.25L Oracle Dicamba Agricultural Herbicide
<i>Oracle Dicamba Herbicide alone controls:</i> Perennial sow thistle Common ragweed English daisy Goldenrod Tansy ragwort Canada thistle Field bindweed Top growth of curled dock False ragweed Giant ragweed	2.3 L Oracle Dicamba Agricultural Herbicide
Diffuse knapweed Goat's beard Ground cherry Pasture sage Poverty weed Sheep sorrel Thyme-leafed spurge	4.6 L Oracle Dicamba Agricultural Herbicide
Baby's breath Fringed sage brush Lambkill Velvet grass <i>And top growth control of:</i> Bracken fern Perennial cinquefoil Russian knapweed	9.2 L Oracle Dicamba Agricultural Herbicide

ROADSIDE VEGETATION CONTROL:

Oracle Dicamba Agricultural Herbicide can be used in tank-mix with Roundup® for annual vegetation control on 1-2 meter wide roadside shoulders. Oracle Dicamba Agricultural Herbicide tank-mixtures with Roundup® and 2,4 D amine offer a broader spectrum of total control of roadside vegetation.

IMPORTANT (Application Directions)

1. Apply 1.25 -2.5 L/ha of Oracle Dicamba Agricultural Herbicide in tank-mix with 0.75-1 L/ha Roundup®.
2. Apply 0.3 L/ha of Oracle Dicamba Agricultural Herbicide in tank-mix with 1.2 L/ha of 2,4-D amine (500g/L formulation) and 0.75 to 1 L/ha of Roundup®.

*Refer to Roundup® label for the appropriate rate of Roundup® for specific weed control and for additional precautions and application instructions.

ESTABLISHED TURF:

Oracle Dicamba Agricultural Herbicide can be used alone or in a tank-mix with 2,4 D amine or 2,4 D L.V. ester herbicide. Oracle Dicamba Agricultural Herbicide tank-mixtures offer additional broadleaf weed control and are particularly effective against dandelion and plantain.

IMPORTANT (Application Directions)

Apply Oracle Dicamba Agricultural Herbicide or Oracle Dicamba Agricultural Herbicide tank-mixtures in at least 110 Litres of water per hectare as a foliar spray to actively growing weeds. Best results will be obtained if application is made in early spring or fall, 2 weeks prior to frost. Apply with caution near trees or shrubs. Do not apply closer than the drip line of trees or shrubs. Do not apply to Bent Grass lawns. Do not rake, mow or water lawn within 24 hours after application. For control

of a broader range of weeds, the recommended rate of Oracle dicamba Agricultural Herbicide may be tank-mixed with 2.2 Litres of 2,4 D amine (500g/L) or 2 Litres of 2,4 D ester (600g/L) herbicide.	
SPECIES CONTROLLED	Rate per Hectare
Clover	1.25 L Oracle Dicamba Agricultural Agricultural Herbicide

Sheep Sorrel

Mouse-eared chickweed

Erect Knotweed

RE-ENTRY INTERVAL:

Do not allow people (other than applicator) or pets on treatment area during application.

Do not enter treated areas until spray has thoroughly dried

CONTAINER CLEANING AND DISPOSAL:

1. Triple-or pressure-rinse the empty container. Add the rinsings to the spray mixture in the tank.
2. Follow provincial instructions for any required additional cleaning of the container prior to its disposal.
3. Make the empty container unsuitable for further use.
4. Dispose of the container in accordance with provincial requirements.
5. For information on the disposal of unused, unwanted product, contact **manufacturer or the provincial regulatory agency**. Contact the manufacturer and the provincial regulatory **agency** in case of a spill, and for clean-up of spills.

Oracle Dicamba **Agricultural** Herbicide alone or with 2,4-D or MCPA

If you have used Oracle Dicamba **Agricultural** Herbicide alone or Oracle Dicamba **Agricultural** Herbicide in a tank mix with 2,4-D or MCPA, to clean the spray equipment follow these steps:

1. Thoroughly hose down the inside and outside of equipment surfaces while filling the spray tank half-full with water. Flush by operating the sprayer until the system is purged of the rinse water.
2. Fill the tank with water, adding 1 L of household ammonia for every 100 L of water. Operate the spray pump to circulate the ammonia solution through the sprayer solution for 15-20 minutes and discharge a small amount of the ammonia solution through the spray boom and nozzles.
3. Flush the solution out of the spray tank through the boom.
4. Remove the nozzles and screens and flush the system with two tanks full of water.

Oracle Dicamba **Agricultural** Herbicide with other herbicides

To clean spray equipment used to apply Oracle Dicamba **Agricultural** Herbicide as a tank mix with wettable powders (WP), emulsifiable concentrates (EC) or other types of water-dispersible formulations, follow these steps: (Note that if you use Oracle Dicamba **Agricultural** Herbicide tank mixes with water-dispersible formulation, you must add detergent to the rinse water.)

1. Thoroughly hose down the inside and outside of equipment surfaces while filling the spray tank half-full with water. Flush by operating the sprayer until the system is purged of the rinse water.
2. Fill tank with water while adding 1 kg of detergent for every 150 litres of water. Operate the pump to circulate the detergent solution through the sprayer system for 5-10 minutes and discharge a small amount of the solution through the boom and nozzles. Let the solution stand for several hours, preferably overnight.
3. Flush the detergent solution out of the spray tank through the boom.
4. Repeat step 1 and follow steps 2 and 3.

Bulk container refilling:

1. The container is to be refilled by the distributor/dealer only with Oracle Dicamba **Agricultural** Herbicide. Do not reuse this container for any other purpose.
2. Reseal and return to an authorized Bulk Site.
3. Prior to refilling, inspect thoroughly for damage such as cracks, punctures, bulges, dents, abrasions and damaged or worn threads on closure devices.
4. Check for leaks after refilling and before transportation.
5. Do not refill or transport damaged or leaking containers.
6. **For information on disposal of unused, unwanted product, contact the manufacturer or the provincial regulatory agency.** Contact the manufacturer and the provincial regulatory **agency** in case of a spill, and for clean-up of spills.
7. If the container is not being refilled, return to the point of purchase (distributor/dealer) for disposal.

STORAGE:

1. Store Oracle Dicamba **Agricultural** Herbicide in its original container only, away from other pesticides, fertilizer, food, or feed.
2. Keep the container closed to prevent spills and contamination.
3. Keep packages dry at all times.

NOTICE TO USER:

This control product is to be used only in accordance with the directions on this label. It is an offence under the *PEST CONTROL PRODUCTS ACT* to use this product in a way that is inconsistent with the directions on the label. The user assumes the risk to persons or property that arises from any such use of this product.

Oracle® Dicamba **Agricultural** Herbicide is a registered trademark of Gharda Chemicals Ltd.

® All other herbicides listed are registered trademarks of their respective companies.